

## Optimization of Ggene Expression and Purification of *Legionella Pneumophila* Peptidoglycan Associated Lipoprotein (PAL) Recombinant Protein

Abolfazl Gholipour<sup>1</sup>; Mojtaba Moosavian<sup>2</sup>; Hamid Galehdari<sup>3</sup>; Manoochehr Makvandi<sup>2</sup>; Ali Mard<sup>4</sup>; Hamid Rajabi memary<sup>5</sup>; Reza Imani<sup>6</sup>; Neda Soleimani<sup>7</sup>; Amirhoushang Alvandi\*<sup>8</sup>

1- Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

2- Department of Microbiology, School of Medicine and Infectious and Tropical Disease Research Center, Ahvaz Jundishapur University of Medical Science, Ahvaz, Iran

3- Department of Genetics, Faculty of Science, Shahid Chamran University, Ahvaz, Iran

4- Department of Physiology, School of Medicine, Ahvaz Jundishapur University of Medical Science, Ahvaz, Iran

5-Department of Agronomy and Plant Breeding, Faculty of Agriculture Shahid University, Ahvaz, IR

6- Molecular and Cellular Research Center, Faculty of Medicine, Shahrekord, Iran

7- Department of Pathology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

8-Department of Medical Microbiology, Faculty of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

gholipour\_abolfazl@yahoo.com

**Background & Objectives:** Legionellae including *Legionella pneumophila*, are recognized as the causative agents for community, travel and hospital-acquired pneumonia. There are many problems with most of available diagnostic tests used to diagnose *Legionella pneumonia*, including inadequate sensitivity and specificity, and inability to provide a result in a clinically useful time period but among them, the urinary antigen tests have revolutionized the laboratory diagnosis of *Legionella pneumonia*. Since the peptidoglycan-associated lipoprotein (PAL) protein of *L. pneumophila* is an urinary antigen and considered as useful diagnostic antigen to diagnose Legionella infection, the aim of this study was to optimaize expression and purification of *L. pneumophila* PAL protein.

**Methods:** In this experimental study, optimizing of 5 parameters (cell density, induction time, growth temperature, IPTG concentration and type of medium) was performed. After expression, periplasmic extract was prepared and recombinant PAL protein purified using Ni<sup>2+</sup>-charged resin column. Finally, recombinant PAL protein was verified by Western blotting.

**Results:** The optimum expression of the r-PAL protein was occurred at an OD<sub>600</sub> of 0.6, 1mM IPTG concentration, after 15 hours at 25°C and use of Terrific Broth medium. Recombinant periplasmic PAL protein was highly purified (>80%) using Ni-NTA column. Also, western blotting analysis showed that recombinant PAL protein was specifically recognized by anti-His6-peroxidase antibody.

**Conclusion:** In order to increase the functional expression of recombinant proteins, a number of strategies were employed such as optimization of expression conditions and periplasmic expression. Also, the results obtained from this study supported the idea that *E. coli* BL21

strain can be served as a functional host for pET26b vector and the designed host and vector can be used in large scale production of recombinant PAL protein

**Keywords:** Optimization; Expression; Purification; Peptidoglycan Associated Lipoprotein (PAL); *Legionella Pneumophila*

